





Effect of exercise on depression and anxiety symptoms: systematic umbrella review with meta-meta-analysis

Neil Richard Munro ¹, Samantha Teague ^{2,3}, Klaire Somoray ²,
Aaron Simpson ^{4,5}, Timothy Budden ⁶, Ben Jackson,⁷ Amanda Rebar,⁸
James Dimmock ^{2,5}

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bjsports-2025-110301>).

¹James Cook University, Townsville City, Queensland, Australia

²Department of Psychology, James Cook University, Townsville City, Queensland, Australia

³Deakin University, Melbourne, Victoria, Australia

⁴University of Western Australia, Perth, Western Australia, Australia

⁵Kids Research Institute Australia, Nedlands, Western Australia, Australia

⁶Central Queensland University, Rockhampton, Queensland, Australia

⁷School of Sport Science, Exercise and Health, University of Western Australia, Crawley, Western Australia, Australia

⁸University of South Carolina, Columbia, South Carolina, USA

Correspondence to

Neil Richard Munro;
neil.munro@jcu.edu.au

Accepted 15 October 2025



© Author(s) (or their employer(s)) 2026. No commercial re-use. See rights and permissions. Published by BMJ Group.

To cite: Munro NR, Teague S, Somoray K, et al. *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2025-110301

ABSTRACT

Objective To synthesise meta-analytic outcomes from randomised controlled trials examining exercise effects on depression and anxiety across all population groups, including children and adults with both clinically diagnosed and subclinical symptoms, excluding those with pre-existing chronic physiological conditions.

Design Meta-meta-analysis (Preferred Reporting Items for Overviews of Reviews (PRIOR) framework).

Data sources Five electronic databases were searched for eligible meta-analyses published from inception to 31 July 2025.

Eligibility criteria for selecting studies Meta-analyses of randomised controlled trials examining exercise interventions for the management of depression and anxiety symptoms were included. To avoid contamination effects, meta-analyses exclusively focused on populations in which chronic physiological conditions were excluded. Study selection was undertaken in duplicate by two independent reviewers.

Results 63 studies (81 meta-analyses, 1079 component studies and 79 551 participants) were included. Exercise reduced depression (standardised mean difference (SMD)=−0.61, 95% CI −0.69 to −0.54) and anxiety (SMD)=−0.47, 95% CI −0.59 to −0.36) symptoms, with aerobic exercise demonstrating the most substantial impact on both depression and anxiety symptoms. The greatest benefits by population group for depression were seen in emerging adults aged 18–30 and postnatal women. Greater reductions in depression were associated with exercise in group and supervised settings. Exercise of shorter duration and at lower intensity was most strongly associated with anxiety reduction.

Conclusion and relevance The findings of the study support that exercise based interventions, in all formats and parameters, can help mitigate depression and anxiety symptoms across all population categories. These results can help health professionals provide targeted, cost effective, evidence based support that aligns with individual profiles and preferences.

Trial registration PROSPERO CRD42020210651.

INTRODUCTION

Depression and anxiety disorders represent significant global health challenges, affecting 7–25% of the population worldwide.^{1–3} These conditions extend beyond psychological problems, impacting family and social functioning,^{4–6} physical health^{7 8} and

WHAT IS ALREADY KNOWN?

- ⇒ Exercise is an effective intervention for reducing symptoms of depression and anxiety, with prior research suggesting comparable benefits to psychotherapy and pharmacotherapy, but there is limited uptake of exercise as a first line treatment.
- ⇒ Previous studies have examined different exercise types, intensities, and population subgroups, but variability in study designs has made it challenging to synthesise findings for clinical application.

WHAT ARE THE NEW FINDINGS?

- ⇒ This meta-meta-analysis is the first to comprehensively isolate the effect of exercise on both depression and anxiety across clinically diagnosed and non-clinical populations, including children, emerging adults, older adults and perinatal women.
- ⇒ Exercise was effective across all population groups, with aerobic, group based and supervised formats showing the greatest benefits for depression.
- ⇒ Differences in the impact of exercise intensity, duration and frequency were revealed for depression and anxiety based symptoms

HOW THIS STUDY MAY IMPACT RESEARCH, PRACTICE OR POLICY?

- ⇒ Mental health professionals should prescribe exercise with the same confidence as traditional treatments, recognising that all exercise formats demonstrate positive effects while tailoring programmes to individual profiles and preferences (e.g., group-based/supervised formats show strongest effects for depression; shorter, lower-intensity programmes for anxiety).
- ⇒ Public health guidelines should position exercise as an accessible, evidence-based first-line intervention for mental health, particularly targeting emerging adults and perinatal populations where effects are strongest

carrying substantial economic costs.⁸ Youth populations experience almost twice the rate of depression and anxiety disorders of adults,^{9–11} with a particularly concerning incidence among women.¹²

While traditional treatments such as antidepressants^{13 14} and psychotherapy¹⁵ have become increasingly common, rising prevalence rates for depression and anxiety symptoms¹⁶ suggest that these therapies alone are not sufficient.^{17 18} Evidence consistently demonstrates that exercise represents a promising intervention, with empirical findings supporting its role in developing cognitive and neurobiological pathways that enhance mental health outcomes.^{19 20} Research consistently indicates the positive impact of exercise on quality of life and psychological well-being in both clinically diagnosed^{21 22} and non-clinically diagnosed populations.²³ While evidence for exercise interventions also shows positive effects on anxiety symptoms,^{24–26} challenges remain regarding how to optimise exercise design to improve adherence rates.²⁷

Meta-meta-analyses have indicated the potential for exercise to mitigate depression and anxiety symptoms,^{23 28} while gaps remain in understanding its effectiveness across age ranges and within exercise parameters. Previous meta-meta-analyses have been limited to adult populations^{23 28} or included populations with confounding factors, such as chronic diseases.²⁹ These were also conducted before the development of the Preferred Reporting Items for Overviews of Reviews (PRIOR) framework,³⁰ which provides comprehensive guidance to facilitate transparency for overviews of healthcare intervention reviews.

In this meta-meta-analysis, our aims were to: (1) provide comprehensive estimates of the impact of exercise on depression and anxiety symptoms across all age ranges, including clinical and non-clinical populations and (2) examine moderation effects of exercise parameters (type, duration, frequency, intensity, supervision and group based activity) and population groups, applying the PRIOR methodology.

METHODS

Protocol and registration

This meta-meta-analysis was conducted following a prospectively registered protocol published in the International Prospective Register of Systematic Reviews (PROSPERO). The only deviations from the protocol were to include additional authors. Our methodological approach adheres to the protocol as described in PROSPERO and with the Reporting Guideline for Overviews of Reviews of Healthcare Interventions.³⁰

Search strategy and selection criteria

Five electronic databases were searched: SCOPUS, PsycINFO, CINAHL, OVID Medline and SPORTDiscus. The search was conducted in October 2023 with no restrictions, and repeated in September 2024 to capture more recent literature. A further search was conducted in July 2025 with all five databases, and two additional databases, Embase and Cochrane Library, were included. Manual searches were also performed. Search terms covered constructs related to depression, anxiety, randomised controlled trial designs, meta-analyses and exercise programmes, and were limited to English language peer reviewed journal articles (online supplemental appendix 1).

Search results were exported to the Colandr tool³¹ for duplicate removal. Two independent reviewers (NRM and AS) screened the titles and abstracts of the identified studies. Disagreements were resolved through discussion between the reviewers and the broader author team. The full text assessment was performed in duplicate (NRM and AS), with additional discussion from the supervisory team (JD, KS and ST) and reached 100% agreement for studies to be included in the final data extraction. In cases

where full texts were unavailable or data clarity was required (five instances), study authors were contacted for clarification.

The PICOS (Patient/Population, Intervention, Comparison, Outcome, Study design) framework was used to define the inclusion and exclusion criteria.³² Eligible studies included those with participants of any age who were assessed for depression and anxiety symptoms or diagnosed with depression or anxiety disorders according to DSM-5,³³ ICD-10³⁴ or validated self-reported measures. Studies that explicitly stated the inclusion of individuals with physiological conditions (eg, heart disease, cancers, HIV, or Parkinson's disease) were excluded to avoid potential confounding effects on the relationship between motivation to exercise and mental health.³⁵

Exercise interventions were defined as planned, structured, repetitive and purposeful physical activities to improve physical and mental health.³⁶ All exercise modalities, intensities, frequencies and settings (individual or group) were considered eligible. Studies that combined exercise with psychotherapy, nutrition, pharmacotherapy treatments or manual therapies, and any other non-exercise based interventions were excluded unless the independent effect of exercise could be isolated.

Only meta-analyses of randomised controlled trials (RCTs) comparing exercise with an active control, placebo or no intervention (eg, waitlist) were included. Meta-analyses that incorporated non-randomised studies or cohort designs were excluded. Eligible meta-analyses assessed the impact of exercise on depression or anxiety symptoms using validated self-report or clinician rated measures. Only meta-analyses reporting standardised mean difference (SMD)³⁷ or Hedges g ³⁸ were eligible. Eligible studies were systematic reviews that included meta-analyses of RCTs meeting the PICO criteria. Narrative, umbrella and systematic reviews without a meta-analysis were excluded.

Data extraction and quality assessment

Data extraction was performed using a systematic template and subjected to further independent audits of potential errors by two additional researchers (KS and AR) to reach 100% verification. The following data were collected: (1) demographic information (age range, sample size, gender distribution and participant characteristics); (2) intervention characteristics (exercise category, intensity, duration, frequency, group or individual exercise format and specific sports); (3) RCT and control group features; (4) depression or anxiety characteristics and assessment scales used; (5) effect scores on depression or anxiety symptoms as SMD or Hedges' g , confidence intervals (CIs), p values, heterogeneity measures (I^2 statistic³⁹) and standard errors; and (6) meta-analysis study characteristics, including certainty of evidence using assessment⁴⁰ and risk of bias analyses. For clarity in reporting results, we state the number of meta-analyses collected for each subsection, although some reviews may include more than one meta-analysis.

The primary author (NRM) assessed the risk of bias in the included systematic reviews and reviewed with the supervisory panel (JD, ST and KS) to reach complete agreement on AMSTAR-2 categorisation.⁴¹ Each included meta-analysis was classified as high, moderate, low or critically low quality (online supplemental table 2). Sixteen items are considered within AMSTAR-2, each scoring as yes, partial or no. Seven items are considered critical and 11 non-critical. To assess for publication bias, additional funnel plots were conducted to review asymmetries.

Data synthesis

Data synthesis was conducted using R Studio software⁴² and the Metafor package.⁴³ Random effects meta-analyses were conducted to examine the overall effect of exercise interventions on depression and anxiety symptoms. SMDs^{37 44} with 95% CIs were used as the effect size measure, interpreted according to Cohen's *d* conventions.⁴⁵ Heterogeneity was assessed using I^2 , with values of 25%, 50% and 75% indicating low, moderate and high heterogeneity, respectively.

To evaluate the robustness of findings, sensitivity analyses were conducted to compare the synthesis of overall depression and anxiety outcomes across studies categorised by AMSTAR⁴¹ ratings (critically low, low, moderate and high). Subgroup analyses were performed to explore differences by population category (youth aged <18 years, emerging adults aged 18–30 years, adults aged ≥18 years and adults aged ≥55 years, and women in prenatal, postnatal and perinatal stages), clinical depression or anxiety status and exercise modality, provided at least two studies reported the effect estimates for the same population, intervention and outcome.

The corrected covered area (CCA) method was used to evaluate the degree of overlap in component studies across all reviews.⁴⁶ This approach quantifies the extent of primary research duplication among systematic reviews, with a CCA of 0% indicating complete uniqueness of component studies across meta-analyses and 100% signifying complete duplication. CCA calculations are interpreted as slight (0–5%), moderate (6–10%), high (11–15%) or very high (>15%). The evidence classification system developed by the Oxford Centre for Evidence Based Medicine (GRADE)⁴⁷ was used to provide additional certainty of evidence.

Equity, diversity and inclusion statement

Our research and author team included three women and five men, consisting of junior, mid-career and senior researchers from different countries. Four of the team members were initially from Australia, two from the UK, one from the US and one from the Philippines. As a meta-meta-analysis, the populations within the component studies span across countries globally. From a financial and time resourcing perspective, we sourced the contributing meta-analyses to those published in English and, in turn, recognise the potential limitation in the applicability of findings to non-English speaking populations.

RESULTS

The study selection process and rationale are outlined in the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) flowchart⁴⁸ (figure 1). The initial database search gave 2517 records, of which 57 met the eligibility criteria.

Characteristics of included reviews

Online supplemental table 1 presents the characteristics of all of the included reviews. The selected meta-analyses for depression (n=57) included 800 component studies, with a total of 57 930 participants aged 10–90 years, diagnosed with clinical levels of depression or experiencing depressive symptoms. Exercise interventions for depression analyses were categorised as aerobic (eg, running, walking and cycling; n=19), with 181 component studies of 9941 participants, resistance (eg, strength training; n=8) with 93 component studies of 4770 participants, mind-body (eg, yoga, tai-chi and qigong; n=16) with 90 component studies of 7257 participants or mixed exercise modalities (n=39), with 631 component reviews of 48 696 participants. For




Figure 1 Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) flowchart of study selection process. NA, not available; RCT, randomised controlled trial.

anxiety, the selected meta-analyses (n=24) comprised 258 component studies of 19 368 participants aged 18–67 years. Exercise interventions for anxiety analyses were categorised as aerobic (n=7), with 32 component studies of 1235 participants, resistance (n=1), with five component reviews of 300 participants, mind–body (n=9), with 74 component studies of 5175 participants or mixed exercise modalities (n=13), with 170 component studies of 14 208 participants.

The overall CCA rating was 1.16% for depression reviews and 0.52% for anxiety reviews, indicating slight overlap. The characteristics of all of the depression reviews are shown in online supplemental table 3, with anxiety reviews in online supplemental table 4. Most reviews (n=34) received a critically low AMSTAR-2 score, while the remainder were distributed between low rated (n=21) and high rated (n=8) reviews (online supplemental table 2). A sensitivity analysis revealed minimal inter-group differences for both depression (online supplemental table 3) and anxiety (online supplemental table 4) by AMSTAR-2 quality level. Using the GRADE framework,⁴⁷ all included reviews were classified as level 1 (meta-analyses of RCTs) and graded as high quality.

Analysis of funnel plots for the incorporated reviews revealed no discernible evidence of publication bias due to the overall symmetrical appearance of the plot for reviews that examined depression. However, a slight sign of asymmetry was observed for the reviews that examined anxiety, as indicated by an imbalance in the distribution of reviews, with more reviews on the left side (adverse effects) than the right side (positive effects) (figure 2a, b).

Depression

Overall analysis

The overall analysis of 50 reviews (57 meta-analyses, 800 component studies and 57 930 participants) showed a significant medium sized SMD^{37 44} of -0.61 (95% CI -0.69 to -0.54).

Subgroup analyses

Exercise was effective in reducing depression symptoms across all population subgroups. Exercise showed the most pronounced effect for emerging adults, represented by four reviews^{49–52} (four meta-analyses, 54 component studies and 4180 participants) with SMD of -0.81 (95% CI -1.06 to -0.57). Exercise interventions across all age groups were effective: SMD -0.66 (95% CI -0.79 to -0.54) for adults, from 23 reviews^{21 53–74} (27 meta-analyses, 428 component studies and 24 154 participants); SMD -0.53 (95% CI -0.81 to -0.24) for youths aged <18 years analysed in six reviews^{75–80} (six meta-analyses, 79 component studies and 7474 participants); SMD -0.81 (95% CI -1.06 to -0.57) for emerging adults aged 18–30 analysed in four reviews^{49 50 81 82} (four meta-analyses, 54 component studies and 4180 participants); and SMD -0.41 (95% CI -0.51 to -0.30) for late adulthood, represented by six reviews^{66 83–87} (eight meta-analyses, 72 component studies and 5860 participants).

Postnatal populations, represented by four reviews^{88–91} (four meta-analyses, 42 component studies and 3437 participants) showed the most pronounced effect of exercise on depression within perinatal groups, with an SMD -0.70 (95% CI -0.92 to -0.48). Exercise was also effective for prenatal populations with SMD -0.46 (95% CI -0.59 to -0.32), analysed in four reviews^{92–95} (four meta-analyses, 42 component studies and 4717 participants) and for the broader perinatal category SMD of -0.52 (95% CI -0.66 to -0.39) analysed in 11 reviews^{88–98} (11 meta-analyses, 150 component studies and 15 518 participants).

The perinatal group incorporated prenatal and postnatal populations and reviews explicitly designated as perinatal. These three categories showed high to very high interpretations in the CCA analysis: postnatal (26.92%), prenatal (18.67%) and perinatal (12.76%).

Exercise modes for depression

Aerobic exercise showed the most substantial impact, with SMD of -0.81 (95% CI -1.01 to -0.60) based on 15 reviews^{52 55 63 65–67 70 74 75 79 80 85 91 97 99} (19 meta-analyses, 181 component studies and 9941 participants). All other exercise modes were effective: -0.62 (95% CI -0.93 to -0.31) for resistance training derived from eight reviews^{52 55 65 67 70 71 74 80 85} (12 meta-analyses, 93 component studies and 4770 participants); -0.53 (95% CI -0.66 to -0.39) for mind–body interventions, from 11 reviews^{51 52 56 60 62 64 69 80 86 87 91 92 97} (16 meta-analyses, 90 component studies and 7257 participants) and -0.60 (95% CI -0.68 to -0.52) for mixed exercise modalities, based on 35 reviews^{21 49 50 52–55 57–59 61 66 68 70 72 73 75–80 83–85 88–91 93–98} (39 meta-analyses, 631 component studies and 48 696 participants).

Group based exercise appeared more effective with SMD of -0.71 (95% CI -0.93 to -0.47) based on seven reviews^{66 68 70 79 80 88 97} (seven meta-analyses, 70 component studies and 4858 participants), compared with individual performed exercise with SMD of -0.38 (95% CI -0.65 to -0.11) reported from five reviews^{66 68 70 79 97} (five meta-analyses, 41 component studies and 2253 participants).

Exercise intensity for depression

From four reviews^{70 72 79 99} (four meta-analyses, 17 component studies and 1063 participants), low intensity exercise showed SMD of -0.69 (95% CI -1.09 to -0.30). Four reviews^{68 70 72 99} for moderate intensity interventions (four meta-analyses, 43 component studies and 3217 participants) showed a more substantial impact with SMD of -1.02 (95% CI -1.68 to -0.35). Moderate to vigorous intensity, from two reviews^{68 79} (two meta-analyses, 11 component studies and 591 participants), showed a comparable effect with SMD of -0.78 (95% CI -1.26 to -0.31). Vigorous intensity, from three reviews^{70 74 99} (three meta-analyses, 17 component studies and 981 participants), showed a comparable effect with SMD of -0.65 (95% CI -0.99 to -0.31).

Exercise duration for depression

Longer term exercise over 24 weeks showed the most substantial impact from two reviews^{69 85} (two meta-analyses, 20 component studies and 987 participants) reporting SMD of -1.11 (95% CI -2.01 to -0.21). Medium term exercise, lasting 9–24 weeks, showed a more modest effect, from four reviews^{52 69 85 86} (seven meta-analyses, 45 component studies and 2874 participants) reporting SMD of -0.44 (95% CI -0.57 to -0.31). Short-term exercise of up to 8 weeks demonstrated a medium effect from four reviews^{50 52 79 95} (four meta-analyses, 21 component studies and 958 participants) with SMD of -0.76 (95% CI: -1.07 to -0.45).

Exercise frequency for depression

Exercise with a higher frequency, ≥ 3 days per week, reported SMD of -0.52 (95% CI -0.75 to -0.29) from four reviews^{50 52 71 79} (six meta-analyses, 39 component studies and 2270 participants), with a similar impact to exercise with lower frequency, occurring 1 or 2 days per week, with SMD of -0.43




Figure 2 Funnel plot analysis of publication bias for anxiety. (A) Funnel plot analysis of publication bias for depression. (B) Funnel plot analysis of publication bias for anxiety.

(95% CI -0.73 to -0.13) from three reviews^{71 81 100} (three meta-analyses, 30 component studies and 1820 participants).

Exercise supervision for depression

Supervised exercise showed a more pronounced impact with SMD of -0.69 (95% CI -1.08 to -0.30) from three reviews^{57 66 70} (three meta-analyses, 93 component studies and 4624 participants) than unsupervised activity SMD of -0.46 (95% CI -0.66 to -0.26) from three reviews^{57 66 70} (three meta-analyses, 35 component studies and 1698 participants).

Exercise for clinical and non-clinical depression

For populations with clinically diagnosed depression, eight reviews^{21 54 58 63 68 75 78 95} (eight meta-analyses, 79 component

studies and 4960 participants) reported SMD of -0.73 (95% CI -0.93 to -0.53) with a slightly larger impact SMD of -0.81 (95% CI -0.98 to -0.64) reported in populations explicitly identified as non-clinically diagnosed from two reviews^{75 78} (two meta-analyses, 24 component studies and 1617 participants).

Anxiety

Overall analysis

The impact of exercise on anxiety symptoms, derived from 23 reviews (24 meta-analyses, 258 component studies and 19 368 participants) showed a small to moderate SMD^{37 44} of -0.47 (95% CI -0.59 to -0.36).

Subgroup analyses

Exercise was effective in reducing depression symptoms across all anxiety subgroups. For emerging adults, four reviews^{50-52 101} (four meta-analyses, 48 component studies and 4186 participants) showed a moderate reduction in anxiety symptoms, with SMD of -0.59 (95% CI -0.65 to -0.53). The adult population, examined in 17 reviews^{56 58 60 62 72 74 102-112} (18 meta-analyses, 184 component studies and 13 186 participants), showed a slightly smaller effect size with SMD of -0.40 (95% CI -0.52 to -0.27).

Exercise modes for anxiety

Aerobic exercise showed the most substantial effect on anxiety symptoms with SMD of -0.60 (95% CI -0.87 to -0.33) from five reviews^{52 74 102 104 107} (five meta-analyses, seven component studies and 1235 participants). All other exercise modes were effective: resistance based exercise SMD of -0.56 (95% CI -0.84 to -0.28) based on one review⁵² (one meta-analysis, five component studies and 300 participants); mind-body exercise SMD of -0.50 (95% CI -0.67 to -0.32) from eight reviews^{56 60 62 81 82 106 108 111} (nine meta-analyses, 74 component studies and 5175 participants); and mixed exercise SMD of -0.45 (95% CI -0.60 to -0.30) from 12 reviews^{50 58 72 81 98 101 103 105 109 110 112 113} (13 meta-analyses, 170 component studies and 14 208 participants). One group based exercise review¹⁰⁴ showed effectiveness with SMD of -0.60 (95% CI -0.94 to -0.26) from one review (one meta-analysis, two component studies and 72 participants), and no meta-analysis data were available for individual performed exercise on anxiety.

Exercise intensity for anxiety

Low intensity exercise showed a more substantial impact with SMD of -0.68 (95% CI -1.21 to -0.14) from two studies^{72 110} (two meta-analyses, nine component studies and 903 participants) compared with the effect of moderate intensity exercise with SMD of -0.06 (95% CI -0.22 to 0.09) reported from the same two studies^{72 110} (two meta-analyses, 13 component studies and 2069 participants). No data were available for moderate to vigorous exercise for anxiety, although data for vigorous exercise showed SMD of -0.17 (95% CI -0.53 to 0.19) from one review⁷⁴ (one meta-analysis, five component studies and 225 participants).

Exercise duration for anxiety

Short term exercise, up to 8 weeks, showed the most substantial impact with SMD of -0.70 (95% CI -0.92 to -0.47) from two reviews^{50 52} (two meta-analyses, 12 component studies and 267 participants) compared with the impact of medium term exercise, between 9 to 24 weeks, with SMD of -0.50 (95% CI -0.64 to -0.37) from two reviews^{52 110} (four meta-analyses, 23 component studies and 2285 participants), and longer term exercise, lasting >24 weeks, with SMD of -0.03 (95% CI -0.18 to 0.13) from one review¹¹⁰ (one meta-analysis, seven component studies and 1013 participants).

Exercise frequency for anxiety

Lower frequency exercise, 1 or 2 days per week, was examined in one review⁵² (one meta-analysis, six component studies and 293 participants) with SMD of -0.71 (95% CI -0.95 to -0.47), which showed more impact than higher frequency exercise of ≥ 3 days per week, with SMD of -0.50 (95% CI -0.71

to -0.30) from two reviews^{50 52} (four meta-analyses, 18 component studies and 1573 participants).

Exercise supervision for anxiety

No meta-analyses were available that reported explicitly on exercise supervision levels for anxiety.

Exercise for clinical and non-clinical anxiety

Two reviews^{107 109} (one meta-analysis, 19 component studies and 935 participants) on clinically diagnosed anxiety were analysed, reporting an SMD of -0.42 (95% CI -0.61 to -0.22). No data were available for the non-clinical anxiety population.

DISCUSSION

This meta-meta-analysis is the first to comprehensively isolate the effect of exercise, by excluding pre-existing physiological conditions, on both depression and anxiety across clinically diagnosed and non-clinical populations, including children, emerging adults, older adults and perinatal women. Exercise had a medium sized effect on depression symptoms and a small-to-medium effect on anxiety symptoms, with the most substantial effects found for emerging adults and perinatal populations, particularly in the postnatal period. All exercise modalities showed positive effects, with aerobic, group based and supervised formats appearing to be the most effective for depression symptoms. Aerobic, resistance, mind-body and mixed exercise modalities showed a medium impact for the mitigation of anxiety symptoms.

Group based and supervised interventions showed notably higher effect sizes than individual based activities for depression, suggesting that social components have a crucial role in the antidepressant effects of exercise. These outcomes align with research demonstrating exercise's social support benefits,¹¹⁴ indicating that the psychological sense of belonging may contribute additional value at both biological and psychosocial levels.¹¹⁵⁻¹¹⁷ Psychological interventions conducted within groups can be more successful than interventions targeted at an individual level, and it is possible that articulating intentions in a group setting may increase motivation to persevere with behaviour change.¹¹⁸ Furthermore, expectations surrounding the impact of exercise can increase when participants elaborate on their expectations, thereby increasing the possibility of rating the intervention as successful.¹¹⁹ Mental health professionals should consider strategies for promoting exercise as a cost effective intervention and examine how contextual factors, particularly social and physical environments, can positively influence outcomes.^{23 120 121}

The effectiveness of exercise interventions appeared comparable with pharmacological treatments (SMD= -0.36) and psychotherapies (SMD= -0.34) for managing depression and anxiety symptoms.¹²² Given the cost effectiveness, accessibility and additional physical health benefits of exercise, these results underscore the potential for exercise as a first line intervention, particularly in settings where traditional mental health treatments may be less accessible or acceptable.^{18 123} Early interventions of exercise have been shown to enhance recovery from physiological conditions,^{124 125} and there is evidence that early interventions, from an age related or diagnosis timing perspective, can support recovery from ICD-10¹²⁶ or DSM-5 diagnosed mental health conditions.¹²⁷ As exercise also stimulates neurobiological mechanisms, including increased growth of brain neurotrophins and protection against neurotoxic damage,^{128 129} our findings further support public health guidance beyond the immediate impact of alleviating depression and anxiety symptoms.

Our findings suggested that exercise interventions showed enhanced effectiveness in ameliorating depressive symptoms when implemented in populations without comorbid physiological conditions, relative to outcomes reported in a previous systematic review examining populations that did not specifically exclude such comorbidities (SMD = -0.43).²⁸ These divergent findings may reflect the impact of pre-existing severe medical conditions on motivation and ability to exercise,^{130 131} alongside potential participation bias for physically healthier populations.¹³² While the overall effect of exercise on all age groups was positive, the analysis of subgroups suggested that there may be nuanced differences in the impact of exercise based on specific population characteristics. For instance, we found particularly strong effects for emerging adults, an age that can typically signify the onset of mental health conditions¹³³ and postnatal women cohorts. That postnatal women may particularly benefit from exercise is especially significant given the high prevalence and potential severity of postpartum depression,¹³⁴ underscoring the potential of exercise as a low risk, high benefit strategy for improving maternal mental health during this vulnerable period.^{135 136}

For anxiety management, lower intensity and shorter duration (up to 8 weeks) interventions seemed more efficacious, although a wide range of definitions was provided for intensity parameters. This pattern suggests that the prescription of more immediate and shorter time frame exercise regimens may offer substantial benefits for individuals with anxiety. Early stage interventions for anxiety disorders have been shown to mitigate the onset of secondary mental health disorders.¹³⁷ Promotion of briefer exercise interventions may, therefore, support populations with anxiety symptoms.

While our study excluded populations affected by factors that might influence exercise engagement and did not explicitly examine motivational factors, it is important to acknowledge that physical and mental health characteristics contribute significantly to exercise adoption.¹³⁸ Incorporating intrinsic motivational factors into exercise programme design can significantly influence intervention adherence and participation rates.^{36 139 140} Therefore, future research on intervention design and exercise prescription should prioritise the consideration of individual circumstances and needs, thereby supporting increased motivation and responses.¹⁴¹

Strengths and limitations of this study

The primary strength of this study was its comprehensive synthesis of data from 72 meta-analyses, incorporating 926 component randomised controlled studies and involving 66 707 participants. Applying the PRIOR framework's complete recommendations enhanced the robustness of these findings.³⁰ Our research distinguished itself by encompassing clinically diagnosed and non-clinical populations across diverse demographics, including children and young people, emerging adults, older adults and perinatal women. To isolate the precise impact of exercise, we excluded meta-analyses that included populations with pre-existing physiological conditions. This methodological decision mitigated the confounding effects of physiological illness related motivations on exercise engagement and its subsequent psychological outcomes.^{35 142}

Inspection of the funnel plots suggested subtle asymmetry in anxiety focused studies, potentially indicating mild publication bias or true effect heterogeneity. Additionally, heterogeneous interpretations of exercise parameters across constituent studies present challenges in reporting precise exercise intensity

and duration boundaries.^{143 144} Analyses of data from studies of depression for perinatal and youth populations gave high to very high CCA⁴⁶ ratings, indicative of substantial duplication among component studies. This finding underscores the necessity for additional, diverse studies focusing on these specific demographic groups. Implementing the CCA method in the analysis facilitated the quantification of component study overlap across meta-analyses, thereby elucidating potential result distortion due to the double counting of duplicate studies. This methodological approach enhances the synthesis of multiple meta-analyses; however, it does not account for the relative sample sizes of individual component studies. Consequently, research with a substantial sample size (eg, 1000 participants) will contribute the same level of input calculation to CCA as a study with a smaller sample size (eg, 10 participants). While the CCA is not deployed to calculate the overall SMD within this meta-meta-analysis, it is imperative to interpret each output in conjunction with the corresponding CCA to contextualise potential levels of component study duplication (as illustrated in online supplemental table 3 and online supplemental table 4).

Despite most meta-analyses being categorised as critically low or low quality according to AMSTAR-2, sensitivity analyses across quality domains showed minimal impact on the overall results. The substantial volume of meta-analyses available for depression studies engendered high confidence in the results across the analysed subgroups.

The heterogeneous interpretations of exercise parameters used within constituent studies presented a significant challenge in reporting exercise intensity and duration results with precise boundaries. Meta-analyses examining exercise intensity used disparate source definitions for intensity, while exercise duration parameters showed wide variations within component studies. This limitation underscores the need for standardised intensity classifications in future investigations to facilitate more precise comparisons.

A notable limitation of this analysis was the need for more anxiety based meta-analytic data for wider ranging population groups, including late adulthood, youth and perinatal populations. This gap in the literature highlights the need for further research to comprehensively understand the effects of exercise on anxiety across the entire lifespan.

Clinical implications

These findings support exercise as a viable and effective intervention for depression and anxiety symptoms across diverse populations. However, clinical adoption of exercise based interventions remains limited.¹⁴⁵ Several factors are crucial for successful implementation, including co-designed approaches and individually tailored interventions.^{146 147} Enhancing clinicians' knowledge base and developing pragmatic guidelines could promote broader adoption of exercise interventions in clinical practice.¹⁴⁸⁻¹⁵⁰

CONCLUSIONS

This meta-meta-analysis provides robust evidence that exercise effectively reduced depression and anxiety symptoms across all age groups, comparable with or exceeding traditional pharmacological or psychological interventions. Group and supervised formats gave the most substantial benefits, underscoring the importance of social factors in mental health interventions. With evidence that different characteristics of exercise appear to impact depression and anxiety at varying magnitudes, tailored exercise programmes must be prescribed. Exercise is

an accessible and cost effective treatment option for depression and anxiety, so these findings must be translated into clear, actionable guidelines for ensuring widespread adoption and long term impact.

Contributors NRM, JD, ST and BJ conceptualised the study, and consulted with AR to build upon the previous meta-meta-analysis conducted in this field. JD was responsible for the acquisition and provision of resources. NRM, JD, ST, KS, AR and AS undertook investigation, methodology and project administration. Data curation involved NRM, KS and AS. NRM, KS, AS and AR directly accessed and verified the underlying data reported in the manuscript. The lead statistician was KS, with NRM, JD, ST, AS and AR also assisting with formal analysis. The manuscript was drafted by NRM, with guidance from JD, KS and ST. JD, ST and KS provided supervision. All authors had full access to all the data in the study, reviewed and edited the manuscript, and had final responsibility for the decision to submit for publication. The guarantor (primary author NRM) accepts full responsibility for the finished work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement statement There has been no patient and public involvement in research from patients or members of the public in the research design, conduct, reporting, or dissemination plans.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. All data collection forms, data extracted for analyses and software scripts are available within the supplementary materials or from the authors on request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Neil Richard Munro <https://orcid.org/0000-0003-1507-569X>

Samantha Teague <https://orcid.org/0000-0002-0487-7307>

Klaire Somoray <https://orcid.org/0000-0001-7521-1425>

Aaron Simpson <https://orcid.org/0000-0002-0660-488X>

Timothy Budden <https://orcid.org/0000-0002-5653-6651>

James Dimmock <https://orcid.org/0000-0002-4475-8169>

REFERENCES

- Weinberger AH, Gbedemah M, Martinez AM, et al. Trends in depression prevalence in the USA from 2005 to 2015: widening disparities in vulnerable groups. *Psychol Med* 2018;48:1308–15.
- Remes O, Brayne C, van der Linde R, et al. A systematic review of reviews on the prevalence of anxiety disorders in adult populations. *Brain Behav* 2016;6:e00497.
- Baxter AJ, Scott KM, Vos T, et al. Global prevalence of anxiety disorders: a systematic review and meta-regression. *Psychol Med* 2013;43:897–910.
- Fekadu W, Mihiretu A, Craig TKJ, et al. Multidimensional impact of severe mental illness on family members: systematic review. *BMJ Open* 2019;9:e032391.
- Simon GE. Social and economic burden of mood disorders. *Biol Psychiatry* 2003;54:208–15.
- van Wijngaarden B, Schene AH, Koeter MWJ. Family caregiving in depression: impact on caregivers' daily life, distress, and help seeking. *J Affect Disord* 2004;81:211–22.
- Firth J, Siddiqi N, Koyanagi A, et al. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry* 2019;6:675–712.
- Hoffman DL, Dukas EM, Wittchen HU. Human and economic burden of generalized anxiety disorder. *Depress Anxiety* 2008;25:72–90.
- Abuse S. Mental health services administration. Key substance use and mental health indicators in the United States: results from the 2016 national survey on drug use and health (HHS publication no. SMA 17-5044. Center for Behavioral Health Statistics and Quality; 2017. Available: <https://www.samhsa.gov/data/sites/default/files/NSDUH-FFR1-2016/NSDUH-FFR1-2016.htm>
- Kessler RC, Chiu WT, Demler O, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005;62:617–27.
- Merikangas KR, He J, Burstein M, et al. Lifetime Prevalence of Mental Disorders in U.S. Adolescents: Results from the National Comorbidity Survey Replication—Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry* 2010;49:980–9.
- Daly M. Prevalence of Depression Among Adolescents in the U.S. From 2009 to 2019: Analysis of Trends by Sex, Race/Ethnicity, and Income. *J Adolesc Health* 2022;70:496–9.
- Kendrick T. Strategies to reduce use of antidepressants. *Br J Clin Pharmacol* 2021;87:23–33.
- Read J, Lewis S, Horowitz M, et al. The need for antidepressant withdrawal support services: Recommendations from 708 patients. *Psychiatry Res* 2023;326:115303.
- Ormel J, Hollon SD, Kessler RC, et al. More treatment but no less depression: The treatment-prevalence paradox. *Clin Psychol Rev* 2022;91:102111.
- Jorm AF, Patten SB, Brugha TS, et al. Has increased provision of treatment reduced the prevalence of common mental disorders? Review of the evidence from four countries. *World Psychiatry* 2017;16:90–9.
- Mulder R, Rucklidge J, Wilkinson S. Why has increased provision of psychiatric treatment not reduced the prevalence of mental disorder? *Aust N Z J Psychiatry* 2017;51:1176–7.
- Wainberg ML, Scorza P, Shultz JM, et al. Challenges and Opportunities in Global Mental Health: a Research-to-Practice Perspective. *Curr Psychiatry Rep* 2017;19:1–10.
- Mikkelsen K, Stojanovska L, Polenakovic M, et al. Exercise and mental health. *Maturitas* 2017;106:48–56.
- Smith PJ, Merwin RM. The Role of Exercise in Management of Mental Health Disorders: An Integrative Review. *Annu Rev Med* 2021;72:45–62.
- Rosenbaum S, Tiedemann A, Sherrington C, et al. Physical activity interventions for people with mental illness: a systematic review and meta-analysis. *J Clin Psychiatry* 2014;75:964–74.
- Granero-Jiménez J, López-Rodríguez MM, Dobarrio-Sanz I, et al. Influence of Physical Exercise on Psychological Well-Being of Young Adults: A Quantitative Study. *Int J Environ Res Public Health* 2022;19:4282.
- Rebar AL, Stanton R, Geard D, et al. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychol Rev* 2015;9:366–78.
- Kandola A, Stubbs B. Exercise and anxiety. *Physical Exercise for Human Health* 2020;2020:345–52.
- Jayakody K, Gunadasa S, Hosker C. Exercise for anxiety disorders: systematic review. *Br J Sports Med* 2014;48:187–96.
- McDowell CP, Dishman RK, Gordon BR, et al. Physical Activity and Anxiety: A Systematic Review and Meta-analysis of Prospective Cohort Studies. *Am J Prev Med* 2019;57:545–56.
- Kandola A, Vancampfort D, Herring M, et al. Moving to Beat Anxiety: Epidemiology and Therapeutic Issues with Physical Activity for Anxiety. *Curr Psychiatry Rep* 2018;20:63:63.
- Singh B, Olds T, Curtis R, et al. Effectiveness of physical activity interventions for improving depression, anxiety and distress: an overview of systematic reviews. *Br J Sports Med* 2023;57:1203–9.
- Collado-Mateo D, Lavín-Pérez AM, Peñacoba C, et al. Key Factors Associated with Adherence to Physical Exercise in Patients with Chronic Diseases and Older Adults: An Umbrella Review. *IJERPH* 2021;18:2023.
- Gates M, Gates A, Pieper D, et al. Reporting guideline for overviews of reviews of healthcare interventions: development of the PRIOR statement. *BMJ* 2022;378:e070849.
- Cheng SH, Augustin C, Bethel A, et al. Using machine learning to advance synthesis and use of conservation and environmental evidence. *Conserv Biol* 2018;32:762–4.
- Spitzer RL, Endicott J, Robins E. Research diagnostic criteria: rationale and reliability. *Arch Gen Psychiatry* 1978;35:773–82.
- American Psychiatric Association D, Association AP. *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, DC: American psychiatric association, 2013.
- WHO. *The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines*. World Health Organization, 1992.
- Busch AM, Ciccolo JT, Puspitasari AJ, et al. Preferences for Exercise as a Treatment for Depression. *Ment Health Phys Act* 2016;10:68–72.
- Teixeira PJ, Carraça EV, Markland D, et al. Exercise, physical activity, and self-determination theory: a systematic review. *Int J Behav Nutr Phys Act* 2012;9:1–30.
- Andrade C. Mean Difference, Standardized Mean Difference (SMD), and Their Use in Meta-Analysis. *J Clin Psychiatry* 2020;81:11349.
- Rosenthal R, Cooper H, Hedges L. Parametric measures of effect size. 1994. Available: <https://psycnet.apa.org/record/1993-99100-015> [Accessed 2 Jun 2021].
- Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- Guyatt GH, Oxman AD, Schünemann HJ, et al. GRADE guidelines: a new series of articles in the Journal of Clinical Epidemiology. *J Clin Epidemiol* 2011;64:380–2.

- 41 Shea BJ, Reeves BC, Wells G, *et al.* AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:4008.
- 42 RStudio. RStudio: integrated development for R. RStudio, PBC, Boston, MA; 2022. Available: <http://www.rstudio.com>
- 43 Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36:1–48.
- 44 Cohen J. *Statistical power analysis for the behavioral sciences*. 2013. Available: <https://www.taylorfrancis.com/books/9781134742707>
- 45 Higgins JPT, Thompson SG, Deeks JJ, *et al.* Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 46 Pieper D, Antoine S-L, Mathes T, *et al.* Systematic review finds overlapping reviews were not mentioned in every other overview. *J Clin Epidemiol* 2014;67:368–75.
- 47 Guyatt G, Oxman AD, Akl EA, *et al.* GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383–94.
- 48 Haddaway NR, Page MJ, Pritchard CC, *et al.* PRISMA2020: An R package and shiny app for producing PRISMA 2020-compliant flow diagrams, with interactivity for optimised digital transparency and open synthesis. *Campbell Systematic Reviews* 2022;18:e1230.
- 49 Guo S, Liu F, Shen J, *et al.* Comparative efficacy of seven exercise interventions for symptoms of depression in college students. *Medicine (Baltimore)* 2020;99:e23058.
- 50 Luo Q, Zhang P, Liu Y, *et al.* Intervention of Physical Activity for University Students with Anxiety and Depression during the COVID-19 Pandemic Prevention and Control Period: A Systematic Review and Meta-Analysis. *IJERPH* 2022;19:15338.
- 51 Lin J, Gao Y, Guo Y, *et al.* Effects of qigong exercise on the physical and mental health of college students: a systematic review and Meta-analysis. *BMC Complement Med Ther* 2022;22:287.
- 52 Huang X, Wang Y, Zhang H. Effects of physical exercise intervention on depressive and anxious moods of college students: A meta-analysis of randomized controlled trials. *Asian J Sport Exerc Psychol* 2023;3:206–21.
- 53 Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomised controlled trials. *BMJ* 2001;322:763.
- 54 Schuch FB, Vancampfort D, Richards J, *et al.* Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *J Psychiatr Res* 2016;77:42–51.
- 55 MeadGE, MorleyW, CampbellP. Exercise for depression. *Cochrane Database Syst Rev* 2008.
- 56 Cramer H, Lauche R, Langhorst J, *et al.* Yoga for depression: a systematic review and meta-analysis. *Depress Anxiety* 2013;30:1068–83.
- 57 Conn VS. Depressive symptom outcomes of physical activity interventions: meta-analysis findings. *Ann Behav Med* 2010;39:128–38.
- 58 Rosenbaum S, Vancampfort D, Steel Z, *et al.* Physical activity in the treatment of post-traumatic stress disorder: A systematic review and meta-analysis. *Psychiatry Res* 2015;230:130–6.
- 59 Morres ID, Hatzigeorgiadis A, Stathi A, *et al.* Aerobic exercise for adult patients with major depressive disorder in mental health services: A systematic review and meta-analysis. *Depress Anxiety* 2019;36:39–53.
- 60 Cramer H, Lauche R, Anheyer D, *et al.* Yoga for anxiety: A systematic review and meta-analysis of randomized controlled trials. *Depress Anxiety* 2018;35:830–43.
- 61 Krogh J, Hjorthøj C, Speyer H, *et al.* Exercise for patients with major depression: a systematic review with meta-analysis and trial sequential analysis. *BMJ Open* 2017;7:e014820.
- 62 Vollbehr NK, Bartels-Velthuis AA, Nauta MH, *et al.* Hatha yoga for acute, chronic and/or treatment-resistant mood and anxiety disorders: A systematic review and meta-analysis. *PLoS One* 2018;13:e0204925.
- 63 Davis ME, McGrane N. Exercise as a treatment for clinical depression in adults and progressing the physiotherapists role: A systematic review and meta-analysis. *PPR* 2021;42:105–18.
- 64 Sani NA, Yusoff SSM, Norhayati MN, *et al.* Tai Chi Exercise for Mental and Physical Well-Being in Patients with Depressive Symptoms: A Systematic Review and Meta-Analysis. *IJERPH* 2023;20:2828.
- 65 Jazayeri D, Zanjani Z, Joekar S, *et al.* Comparison Study of the Effects of Resistance and Endurance Exercise Interventions for the Treatment of Depression: A Meta-Analytical Review. *J Contemp Psychother* 2022;52:237–46.
- 66 Lee J, Gierc M, Vila-Rodriguez F, *et al.* Efficacy of exercise combined with standard treatment for depression compared to standard treatment alone: A systematic review and meta-analysis of randomized controlled trials. *J Affect Disord* 2021;295:1494–511.
- 67 Nebiker L, Lichtenstein E, Minghetti A, *et al.* Moderating Effects of Exercise Duration and Intensity in Neuromuscular vs. Endurance Exercise Interventions for the Treatment of Depression: A Meta-Analytical Review. *Front Psychiatry* 2018;9:305:305.
- 68 Correia ÉM, Monteiro D, Bento T, *et al.* Analysis of the Effect of Different Physical Exercise Protocols on Depression in Adults: Systematic Review and Meta-analysis of Randomized Controlled Trials. *Sports Health: A Multidisciplinary Approach* 2024;16:285–94.
- 69 Zeng L, Zhao X, Yu Y, *et al.* Effects of Tai Chi on depression of middle-aged and older adults: an updated systematic review and meta-analysis. *BMC Complement Med Ther* 2023;23:382.
- 70 Heissel A, Heinen D, Brokmeier LL, *et al.* Exercise as medicine for depressive symptoms? A systematic review and meta-analysis with meta-regression. *Br J Sports Med* 2023;57:1049–57.
- 71 Rossi FE, dos Santos GG, Rossi PAQ, *et al.* Strength training has antidepressant effects in people with depression or depressive symptoms but no other severe diseases: A systematic review with meta-analysis. *Psychiatry Res* 2024;334:115805.
- 72 Yue H, Yang Y, Xie F, *et al.* Effects of physical activity on depressive and anxiety symptoms of women in the menopausal transition and menopause: a comprehensive systematic review and meta-analysis of randomized controlled trials. *Int J Behav Nutr Phys Act* 2025;22:13–undefined.
- 73 Han B, Duan Y, Zhang P, *et al.* Effects of exercise on depression and anxiety in postmenopausal women: a pairwise and network meta-analysis of randomized controlled trials. *BMC Public Health* 2024;24:1816.
- 74 Gaia JWP, Schuch FB, Ferreira RW, *et al.* Effects of high-intensity interval training on depressive and anxiety symptoms in healthy individuals: A systematic review and meta-analysis of randomized clinical trials. *Scand Med Sci Sports* 2024;34.
- 75 Bailey AP, Hetrick SE, Rosenbaum S, *et al.* Treating depression with physical activity in adolescents and young adults: a systematic review and meta-analysis of randomised controlled trials. *Psychol Med* 2018;48:1068–83.
- 76 Carter T, Morres ID, Meade O, *et al.* The Effect of Exercise on Depressive Symptoms in Adolescents: A Systematic Review and Meta-Analysis. *J Am Acad Child Adolesc Psychiatry* 2016;55:580–90.
- 77 Axelsdóttir B, Biedilae S, Sagatun Å, *et al.* Review: Exercise for depression in children and adolescents – a systematic review and meta-analysis. *Child Adolesc Ment Health* 2021;26:347–56.
- 78 Wang X, Cai Z-D, Jiang W-T, *et al.* Systematic review and meta-analysis of the effects of exercise on depression in adolescents. *Child Adolesc Psychiatry Ment Health* 2022;16:16.
- 79 Zhang CS, Cheng L, Chen X, *et al.* The strategies of exercise intervention for adolescent depression: A meta-analysis of randomized controlled trials. *Front Psychol* 2022;13:974382.
- 80 Li J, Zhou X, Huang Z, *et al.* Effect of exercise intervention on depression in children and adolescents: a systematic review and network meta-analysis. *BMC Public Health* 2023;23.
- 81 Huang X, Wang Y, Zhang H. Effects of physical exercise intervention on depressive and anxious moods of college students: A meta-analysis of randomized controlled trials. *AJSEP* 2023;3:206–21.
- 82 Lin J, Gao Y, Guo Y, *et al.* Effects of qigong exercise on the physical and mental health of college students: a systematic review and meta-analysis. *BMC Complement Med Ther* 2022;22.
- 83 Bridle C, Spanjers K, Patel S, *et al.* Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. *Br J Psychiatry* 2012;201:180–5.
- 84 Park S-H, Han KS, Kang C-B. Effects of exercise programs on depressive symptoms, quality of life, and self-esteem in older people: a systematic review of randomized controlled trials. *Appl Nurs Res* 2014;27:219–26.
- 85 Mahmoudi A, Amirshaghaghghi F, Aminzadeh R, *et al.* Effect of Aerobic, Resistance, and Combined Exercise Training on Depressive Symptoms, Quality of Life, and Muscle Strength in Healthy Older Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Biol Res Nurs* 2022;24:541–59.
- 86 Chi I, Jordan-Marsh M, Guo M, *et al.* Tai chi and reduction of depressive symptoms for older adults: A meta-analysis of randomized trials. *Geriatr Gerontol Int* 2013;13:3–12.
- 87 Ko K, Kwok ZCM, Chan HY. Effects of yoga on physical and psychological health among community-dwelling older adults: A systematic review and meta-analysis. *Int J Older People Nursing* 2023;18:e12562.
- 88 Pritchett RV, Daley AJ, Jolly K. Does aerobic exercise reduce postpartum depressive symptoms? a systematic review and meta-analysis. *Br J Gen Pract* 2017;67:e684–91.
- 89 Daley A, Jolly K, MacArthur C. The effectiveness of exercise in the management of post-natal depression: systematic review and meta-analysis. *Fam Pract* 2009;26:154–62.
- 90 Carter T, Bastounis A, Guo B, *et al.* The effectiveness of exercise-based interventions for preventing or treating postpartum depression: a systematic review and meta-analysis. *Arch Womens Ment Health* 2019;22:37–53.
- 91 Wang J, Carru C, Sedda S, *et al.* Comparative impact of exercise-based interventions for postpartum depression: A Bayesian network meta-analysis. *Int J Gynecol Obstet* 2024;165:67–75.
- 92 Gong H, Ni C, Shen X, *et al.* Yoga for prenatal depression: a systematic review and meta-analysis. *BMC Psychiatry* 2015;15:1–8.
- 93 Daley A, Foster L, Long G, *et al.* The effectiveness of exercise for the prevention and treatment of antenatal depression: systematic review with meta-analysis. *BJOG* 2015;122:57–62.

- 94 Sánchez-Polán M, Franco E, Silva-José C, *et al.* Exercise During Pregnancy and Prenatal Depression: A Systematic Review and Meta-Analysis. *Front Physiol* 2021;12:640024.
- 95 Liu L, Liu C, Liu X, *et al.* Summary of the effect of an exercise intervention on antenatal depression and the optimal program: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2023;23:293.
- 96 Morres ID, Tzouma N-A, Hatzigeorgiadis A, *et al.* Exercise for perinatal depressive symptoms: A systematic review and meta-analysis of randomized controlled trials in perinatal health services. *J Affect Disord* 2022;298:26–42.
- 97 Liu X, Wang G, Cao Y. Physical exercise interventions for perinatal depression symptoms in women: A systematic review and meta-analysis. *Front Psychol* 2022;13:1022402.
- 98 Yu H, Mu Q, Lv X, *et al.* Effects of an exercise intervention on maternal depression, anxiety, and fatigue: a systematic review and meta-analysis. *Front Psychol* 2024;15:1473710.
- 99 Li W, Liu Y, Deng J, *et al.* Influence of aerobic exercise on depression in young people: a meta-analysis. *BMC Psychiatry* 2024;24:571.
- 100 Zhang CS, Cheng L, Chen X, *et al.* The strategies of exercise intervention for adolescent depression: A meta-analysis of randomized controlled trials. *Front Psychol* 2023;13.
- 101 Chen P, Mazalan NS, Koh D, *et al.* Effect of exercise intervention on anxiety among college students: a meta-analysis. *Front Psychol* 2025;16:1536295.
- 102 Stubbs B, Vancampfort D, Rosenbaum S, *et al.* An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Res* 2017;249:102–8.
- 103 Ensari I, Greenlee TA, Motl RW, *et al.* Meta-analysis of acute exercise effects on state anxiety: an update of randomized controlled trials over the past 25 years. *Depress Anxiety* 2015;32:624–34.
- 104 Robertson R, Robertson A, Jepson R, *et al.* Walking for depression or depressive symptoms: A systematic review and meta-analysis. *Ment Health Phys Act* 2012;5:66–75.
- 105 Conn VS. Anxiety outcomes after physical activity interventions: meta-analysis findings. *Nurs Res* 2010;59:224–31.
- 106 Wang F, Lee E-KO, Wu T, *et al.* The effects of tai chi on depression, anxiety, and psychological well-being: a systematic review and meta-analysis. *Int J Behav Med* 2014;21:605–17.
- 107 Aylett E, Small N, Bower P. Exercise in the treatment of clinical anxiety in general practice - a systematic review and meta-analysis. *BMC Health Serv Res* 2018;18:559:559.
- 108 Gallegos AM, Crean HF, Pigeon WR, *et al.* Meditation and yoga for posttraumatic stress disorder: A meta-analytic review of randomized controlled trials. *Clin Psychol Rev* 2017;58:115–24.
- 109 Ramos-Sanchez CP, Schuch FB, Seedat S, *et al.* The anxiolytic effects of exercise for people with anxiety and related disorders: An update of the available meta-analytic evidence. *Psychiatry Res* 2021;302:114046.
- 110 Martínez-Domínguez SJ, Lajusticia H, Chedraui P, *et al.* The effect of programmed exercise over anxiety symptoms in midlife and older women: a meta-analysis of randomized controlled trials. *Climacteric* 2018;21:123–31.
- 111 Zoogman S, Goldberg SB, Voursora E, *et al.* Effect of yoga-based interventions for anxiety symptoms: A meta-analysis of randomized controlled trials. *Spiritual Clin Pract (Wash D C)* 2019;6:256–78.
- 112 Björkman F, Ekblom Ö. Physical Exercise as Treatment for PTSD: A Systematic Review and Meta-Analysis. *Mil Med* 2022;187:e1103–13.
- 113 Carter T, Pascoe M, Bastounis A, *et al.* The effect of physical activity on anxiety in children and young people: a systematic review and meta-analysis. *J Affect Disord* 2021;285:10–21.
- 114 Burke SM, Carron AV, Eys MA, *et al.* Group versus individual approach? A meta-analysis of the effectiveness of interventions to promote physical activity. *bpspspr* 2006;2:13–29.
- 115 Eys M, Bruner MW, Martin LJ. The dynamic group environment in sport and exercise. *Psychol Sport Exerc* 2019;42:40–7.
- 116 Kandola A, Ashdown-Franks G, Hendrikse J, *et al.* Physical activity and depression: Towards understanding the antidepressant mechanisms of physical activity. *Neurosci Biobehav Rev* 2019;107:525–39.
- 117 Kredlow MA, Capozzoli MC, Hearon BA, *et al.* The effects of physical activity on sleep: a meta-analytic review. *J Behav Med* 2015;38:427–49.
- 118 Steinmetz H, Knapstein M, Ajzen I, *et al.* How Effective are Behavior Change Interventions Based on the Theory of Planned Behavior? *Zeitschrift Für Psychologie* 2016;224:216–33.
- 119 Helfer SG, Elhai JD, Geers AL. Affect and exercise: positive affective expectations can increase post-exercise mood and exercise intentions. *Ann Behav Med* 2015;49:269–79.
- 120 Vella SA, Sutcliffe JT, Fernandez D, *et al.* Context matters: A review of reviews examining the effects of contextual factors in physical activity interventions on mental health and wellbeing. *Ment Health Phys Act* 2023;25:100520.
- 121 Ekkekakis P. Why Is Exercise Underutilized in Clinical Practice Despite Evidence It Is Effective? Lessons in Pragmatism From the Inclusion of Exercise in Guidelines for the Treatment of Depression in the British National Health Service. *Kinesiol Rev (Champaign)* 2020;10:29–50.
- 122 Leichsenring F, Steinert C, Rabung S, *et al.* The efficacy of psychotherapies and pharmacotherapies for mental disorders in adults: an umbrella review and meta-analytic evaluation of recent meta-analyses. *World Psychiatry* 2022;21:133–45.
- 123 Kazdin AE, Rabbitt SM. Novel Models for Delivering Mental Health Services and Reducing the Burdens of Mental Illness. *Clin Psychol Sci* 2013;1:170–91.
- 124 Peixoto TCA, Begot I, Bolzan DW, *et al.* Early exercise-based rehabilitation improves health-related quality of life and functional capacity after acute myocardial infarction: a randomized controlled trial. *Can J Cardiol* 2015;31:308–13.
- 125 Burtin C, Clerckx B, Robbeets C, *et al.* Early exercise in critically ill patients enhances short-term functional recovery. *Crit Care Med* 2009;37:2499–505.
- 126 Gaebel W, Stricker J, Kerst A. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. *Dialogues Clin Neurosci* 2020;22:7–15.
- 127 Association AP. Diagnostic and statistical manual of mental disorders: dsm-5: american psychiatric association. 2013.
- 128 Larsen LQ, Schnor H, Tersbøl BP, *et al.* The impact of exercise training complementary to early intervention in patients with first-episode psychosis: a qualitative sub-study from a randomized controlled feasibility trial. *BMC Psychiatry* 2019;19:1–12.
- 129 Wegner M, Helmich I, Machado S, *et al.* Effects of exercise on anxiety and depression disorders: review of meta-analyses and neurobiological mechanisms. *CNS Neurol Disord Drug Targets* 2014;13:1002–14.
- 130 Murphy C-L, Sheane BJ, Cunnane G. Attitudes towards exercise in patients with chronic disease: the influence of comorbid factors on motivation and ability to exercise. *Postgrad Med J* 2011;87:96–100.
- 131 Tulloch H, Reida R, D'Angelos MS, *et al.* Predicting short and long-term exercise intentions and behaviour in patients with coronary artery disease: A test of protection motivation theory. *Psychol Health* 2009;24:255–69.
- 132 Barreto P de S, Ferrandez A-M, Saliba-Serre B. Are Older Adults Who Volunteer to Participate in an Exercise Study Fitter and Healthier Than Nonvolunteers? The Participation Bias of the Study Population. *J Phys Act Health* 2013;10:359–67.
- 133 McGrath JJ, Al-Hamzawi A, Alonso J, *et al.* Age of onset and cumulative risk of mental disorders: a cross-national analysis of population surveys from 29 countries. *Lancet Psychiatry* 2023;10:668–81.
- 134 Liu X, Wang S, Wang G. Prevalence and Risk Factors of Postpartum Depression in Women: A Systematic Review and Meta-analysis. *J Clin Nurs* 2022;31:2665–77.
- 135 Davenport MH, McCurdy AP, Mottola MF, *et al.* Impact of prenatal exercise on both prenatal and postnatal anxiety and depressive symptoms: a systematic review and meta-analysis. *Br J Sports Med* 2018;52:1376–85.
- 136 Cai C, Busch S, Wang R, *et al.* Physical activity before and during pregnancy and maternal mental health: A systematic review and meta-analysis of observational studies. *J Affect Disord* 2022;309:393–403.
- 137 Kessler RC, Ruscio AM, Shear K, *et al.* Epidemiology of anxiety disorders. *Behavioral Neurobiology of Anxiety and Its Treatment* 2010;2010:21–35.
- 138 Knapen J, Vancampfort D, Moriën Y, *et al.* Exercise therapy improves both mental and physical health in patients with major depression. *Disabil Rehabil* 2015;37:1490–5.
- 139 Deci EL, Ryan RM. Self-determination theory. *Handbook of Theories of Social Psychology* 2012;1:416–36.
- 140 Ryan RM, Patrick H. Self-determination theory and physical 2009. 2009. Available: https://selfdeterminationtheory.org/SDT/documents/2009_RyanWilliamsPatrickDeci_HJOP.pdf
- 141 Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* 2011;6:42.
- 142 Glowacki K, Duncan MJ, Gainforth H, *et al.* Barriers and facilitators to physical activity and exercise among adults with depression: A scoping review. *Ment Health Phys Act* 2017;13:108–19.
- 143 Chekroud SR, Gueorguieva R, Zheutlin AB, *et al.* Association between physical exercise and mental health in 1.2 million individuals in the USA between 2011 and 2015: a cross-sectional study. *Lancet Psychiatry* 2018;5:739–46.
- 144 Garber CE, Blissmer B, Deschenes MR, *et al.* American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43:1334–59.
- 145 Schuch FB, Vancampfort D. Physical activity, exercise, and mental disorders: it is time to move on. *Trends Psychiatry Psychother* 2021;43:177–84.
- 146 Deenik J, Czosnek L, Teasdale SB, *et al.* From impact factors to real impact: translating evidence on lifestyle interventions into routine mental health care. *Transl Behav Med* 2020;10:1070–3.
- 147 Teo JL, Zheng Z, Bird SR. Identifying the factors affecting 'patient engagement' in exercise rehabilitation. *BMC Sports Sci Med Rehabil* 2022;14.
- 148 Garvey L, Benson AC, Bengner D, *et al.* The perceptions of mental health clinicians integrating exercise as an adjunct to routine treatment of depression and anxiety. *Int J Mental Health Nurs* 2023;32:502–12.
- 149 Reed JL, Pipe AL. Practical Approaches to Prescribing Physical Activity and Monitoring Exercise Intensity. *Can J Cardiol* 2016;32:514–22.
- 150 Hill A-M, Hoffmann T, McPhail S, *et al.* Factors Associated With Older Patients' Engagement in Exercise After Hospital Discharge. *Arch Phys Med Rehabil* 2011;92:1395–403.